

Memory formation in the fly brain ショウジョウバエ脳における記憶形成の研究

Dr. Hiromu Tanimoto (谷本 拓 博士)

Head of Independent Junior Research Group Max Planck Institute of Neurobiology, Germany

日時: 2011年12月27日(火)13:30-15:00

会場: 総合研究棟A棟 107 室

Animals can adapt their behaviour to given environment according to memories of their former experiences. To address neural circuits underlying adaptive behaviour, we chose associative memory of Drosophila melanogaster as a model system. Flies form positive or negative memories of an odour by paired presentation of sugar reward or electric shock punishment. Over the past years, we have worked on the neuronal mechanisms for endowing positive or negative values with the odour and found the important roles of neuromodulator dopamine. The value signals by dopamine converge with the odour signal in the mushroom body that consists of second-order olfactory interneurons, thereby the fly forms associative memories. Since dopamine is synthesized in ~280 neurons in the fly brain and involved also in other brain functions, it is important to identify individual responsible neurons for value signalling. I will summarize our most recent findings particularly how we identified different types of neurons for signalling positive and negative values and discuss about the subcellular modulation of synapses at the formation of odour memories.

皆様奮ってご参加下さい。

連絡先: 生命環境科学研究科・若手イニシアティブ・丹羽 隆介 内線 6652 or 4907; rniwa@biol.tsukuba.ac.jp